

Agricultural Production Systems and Climate Change Ductility Enhancement in the Fitri Watershed in Chad

Dona Adoum^{1,*}, Lamy Lamy Georges Maxime², Anguessin Benjamine³, Fawa Guidawa², Mapongmetsem Pierre Marie²

¹Department of Life Sciences and Earth, Faculty of Live Sciences, Earth and Territory planning, University of Sciences and Technology of Ati, Ati, Chad

²Department of Biological Sciences, Faculty of Science, University of Ngaoundere, Ngaoundere, Cameroon

³Department of Environmental Sciences, The Higher National Polytechnic College, University of Maroua, Maroua, Cameroon

Email address:

piermapong@yahoo.fr (Dona Adoum)

*Corresponding author

To cite this article:

Dona Adoum, Lamy Lamy Georges Maxime, Anguessin Benjamine, Fawa Guidawa, Mapongmetsem Pierre Marie. Agricultural Production Systems and Climate Change Ductility Enhancement in the Fitri Watershed in Chad. *American Journal of Agriculture and Forestry*. Vol. 11, No. 4, 2023, pp. 119-127. doi: 10.11648/j.ajaf.20231104.11

Received: April 23, 2023; **Accepted:** May 12, 2023; **Published:** July 6, 2023

Abstract: Chad is one of the most vulnerable countries affected by climate changes. The combination of poverty, demographic pressure and overexploitation of natural resources coupled with the risks of drought and flooding, expose the country to numerous humanitarian emergencies. The objective of this study is to understand the role of agricultural production systems in building resilience to climate change in the Fitri watershed in Chad. The *Faidherbia albida* agroforestry system increases water infiltration and soil fertility rates and conserves biodiversity by limiting the consequences of climate change. Mixed cropping significantly reduces the risk of crop failure due to climatic variations. The survey was carried out on a sample of 178 households divided into 49, 57 and 72 households, respectively in the Mochi, Zegue and Yao areas. The floristic inventory was carried out using 50m square plots. A total of 165 plots were determined in the different farming systems. The area under each agricultural crop varies from one village to another. The ecologically important species are *Faidherbia albida*, *Balanites aegyptiaca*, *Prosopis juliflora* and *Mangifera indica*. Regardless of the area studied, the Fabaceae, Balanitaceae, Mimosaceae, Arecaceae, Rhamnaceae, Capparaceae, Meliaceae and Asclepiadaceae families have an IVF of 10 or more. The high density of woody plants in the agrosystems shows that the individuals are full of seed potential that could contribute to their renewal. The height dominance of the trees implies their carbon sequestration capacity since the sequestered carbon accumulates in the above-ground and below-ground biomass.

Keywords: Agroforestry Systems, Resilience, Climate Change, Biological Diversity, Fitri, Chad

1. Introduction

Chad is one of the vulnerable countries affected by climate changes. The combination of poverty, demographic pressure and overexploitation of natural resources to which drought and flooding risks are added, expose the country to numerous humanitarian urgency and fight efficiently against climate change consequences [1]. Rural populations are the most vulnerable and the most exposed to climate changes. The watershed of the lake Fitri of Chad is a semi-humid ecosystem situated in the Sahelian zone and keep reduces

because of the drought and the increase of water capture for irrigation [2, 3]. This situation impacts crops yield, pastures and population livelihoods. However, the watershed of Fitri is characterized by humid and dry periods of interchanges. The vegetation remains green there for a long part of the year and constitutes a refuge habitat for biodiversity [4]. Soil fertility, water presence and vegetal cover are in proportion favourable factors to activities of diversification inside the Chadian watershed [5]. These have permitted the development of many activities among which agriculture, breeding and fishery according to years and seasons. The

diversity of trees conserved in production systems by farmers varies in function of social, cultural, ecological and economic order needs [6, 7]. The establishment and maintenance of the ecological equilibrium in a sustainable way in Fitri pass through rational management of natural resources [8]. The necessity to improve natural resources management sustainably refers to agroforestry practices [9, 10]. These practices are land managements which associate woods and annual crops and/or animals in spatial conciliation and in time [11]. For many years, agroforestry raises interest in farmers of the Fitri zone of Chad. It is among the alternatives which can mitigate the actual trend which damages natural ecosystems. It focus on the valorization of soil management practices following the environment in its agroecological, economical and cultural context and permits to increase in total production of the cultivated systems [12]. Agroforestry is made up of multifunctional agriculture. It addresses questions of soil fertility loss, rehabilitation of degraded soil, restoration of biodiversity and sequestration of carbon [13]. However, agroforestry is threatened by climate change effects such as desertification, irregularity of rain, high temperature and floods. Among agroforestry species targeted to mitigate these effects of climate changes in the Fitri watershed of Chad, appears *Faidherbia albida*. This species plays a fundamental role in agropastoral activities mainly in soil fertility and forage for livestock. The purpose of the present study is to understand the function of agricultural production systems in the enhancement of ductility against climate changes in the Fitri watershed of Chad. Specifically, it is to: determine farmer strategies developed by farmers to mitigate climate changes; evaluate the exploitation size of different lands; identify wood species suitable to palliate climate changes and determine floristic parameters of the different production systems.

2. Materials and Methods

2.1. Study Site

The study was made in three villages (Yao, Mochi and Zegue), located in Fitri Division in the province of East – Batha of Chad (Table 1). The choice of these lands was based on the presence of agroforestry systems in which farmers practice association of annual crops with trees and/or animals [9, 14].

Table 1. Geographic characteristic data of villages explorés.

Village	Latitudes	Longitude	Altitude (m)
Yao	12° 85' 093"	17° 55' 9.24 "	280
Mochi	12° 80 ' 604"	17° 60' 8.35"	290
Zegue	13° 03 ' 627"	17° 41' 8.48"	300

They are located in Sahelian zones with thermic schemes ranging from 23 to 44°C. They present important constraints between the Northern arid part and the South sahelo-soudanian characterized by a rainfall oscillating between 500 and 800 mm [15]. These contrasts determine different activities between the North with pastoral vocation and the

South where the rainfall conditions permits agricultural practices associated with sedentary semi-transhumant breeding [16]. The uneven is a little damaged and covered by Endoreique Rivers the most important is Lake Fitri where the duration of flow depends on rainfall variability. Five types of soil are distinguished: vertisols, steppe soils, holomorphic soils, hydromorphic soils and mineral raw soils [17]. Three occupation zones are distinguished: pasture zone, agricultural and mixed zones [4]. The vegetation corresponds to the forest with dense wood of different plant groups: forest of *Acacia nilotica* at the border of the lake, gallery forest at the edge of rivers and forest of *Hyphaene thebaica* located in the northwest of the lake [18]. The savannah regroups a woody formation and perennial herbaceous stratus [5]. The lake of Fitri zone is an excellent refuge medium which assures by its multiple resources (water, fish, pasture, wood) a vital role for neighbouring populations. Nevertheless, it is particularly sensitive to climate change effects.

2.2. Methodological Approach

2.2.1. Household Surveys

The methodology of inquiry deals with semi-structural interviews with a first elaborate questionnaire. Farmers were the head of the household and were chosen randomly. A total of 178 households participated in interviews of which 49, 57 and 72 were respectively in Mochi, Zegue and Yao. These numbers were done in function of the census of each village [19, 20]. The main rubric of the questionnaire was based on agricultural production systems, farmer perception of climate changes and endogenous strategies to reinforce their resilience.

2.2.2. Floristic Inventory

Agricultural production systems identified in the watershed of Fitri in Chad are *Faidherbia albida* agroforestry parklands, mixed crops systems and farms of sorghum established out individually (*Sorghum durra*). In each of these systems, plots of 50 m x 50 m were delimited. Inside each of them, trees, shrubs and lianas of diameter at breast height (DBH) or more than 1.30m were measured with a double decimeter and the height of trees was determined with the calliper [21]. A total of 165 plots were registered and distributed as follows: 69, 39 and 57 in Yao, Zegue and Mochi respectively. The geographic data of each site was recorded with a GPS (Global Positioning System).

2.2.3. Data Analysis

Different floristic parameters of the agricultural production systems of Fitri watershed were determined. The Importance Value Index (IVI) of species [22] was used:

$$IVI = 100 * \left[\left(\frac{n_i}{N} \right) + \left(\frac{g_i}{G} \right) + \left(\frac{f_i}{F} \right) \right]$$

with $\frac{n_i}{N} \times 100$ = relative density of the species i; $\frac{g_i}{G}$ = the basal area of the i specie; $\frac{f_i}{F} \times 100$ = The relative

frequency of the species i .

This index permits the evaluation of the specific preponderance of the plant population. Each species with the importance value index is equal to or more than 10% ($IVI \geq 10\%$) is considered ecologically important [23, 24]. In the same way, the Family Importance Index (FIV) [25], was equally used. Each family in which the FIV is equal to or more than 10% ($FIV \geq 10\%$) is ecologically important [26-28]:

$$FIV = 100 * \left[\frac{N_i}{\sum N_i} + \frac{G_i}{\sum G_i} + \frac{D_r}{100} \right]$$

N_i = the number of species of the family i ; G_i = basal area of the species in the family i ; D_r = relative density or the number of species in the family i .

The allometric equation developed in Senegal [29] was successfully applied in similar ecosystems [13] was used to calculate the biomass of woody species.

$$BM \text{ (kg)} = 25.58h - 7.006h^2 + 0.662h^3$$

BM = Biomass (t); h = the height (m). The carbon is determined by the following conversion [30, 31]:

$$C = 0.47 * BM$$

C = Carbon (tC/ha) and BM (t/ha). The carbon stock (tC/ha) is converted into CO_2 equivalent absorbed using the ratio 44/12 (=3.67) corresponding to the ratio CO_2/C [32].

The density and mean diameter of woody species of each production system was calculated according to the following formula.

$$D = \text{Density (Stems/ha)} = n/S * 100$$

n = number of individuals in the plot;

S = area of the plot (ha).

$$DM = \sqrt{\frac{1}{n} \sum_{i=1}^n d_i^2}$$

DM represents the mean diameter in cm, n is the number of stems measured, d_i is the diameter at the breast height of the stem i at 1.30 m above ground.

Data collected were subjected to an analysis of variance (ANOVA). Significant means were separated by the Duncan Multiple Range Test. The statistical program used for the ANOVA was Statgraphics plus 5.0. The Excel spreadsheet was used to draw the graphs. The program Microsoft Excel 2016 serves to classify numeric data and draw graphs.

3. Results

3.1. Farmer Perceptions on Climate Changes in Fitri Watershed

The opinion of farmers is diversified concerning climate change perceptions. The climate parameters mostly announced are among others temperature amplification

(87.93%), irregularity of rainfall (77.59%) and flood (68.97%) (Table 2).

Table 2. Farmer climate change perceptions.

Perceptions	Farmer answers (%)		
	Yes	No	neutral
Desertification	12.07	68.97	18.97
Temperature growth	87.93	1.2	10.34
Irregular Rainfall	77.59	8.62	13.79
Floods	68.97	22.44	8.62

The majority of farmers (87.93%) interviewed estimate that temperature is the main reason for climate changes whereas 68.97% of them are unanimous that, desertification is not involved in climate changes. The irregularity of rain and high temperatures modify the phenology of crops.

3.2. Endogenous Resilience Strategies Against Climate Changes

Various endogenous strategies were adopted by farmers against climate changes. They developed powerful agricultural production systems among which the key strategy was the *Faidherbia albida* agroforestry parklands. They associate cereal crops with *Faidherbia albida*. This multipurpose tree presents a reverse phenology. Its shades off its leaves during the rainy season and recovers them during the dry one [33, 34]. Farmers are unanimous that this Fabaceae is an agroecological species. Its presence in the agricultural production system improves soil fertility, protects crops against wind erosion and controls the sand flow. The yield of sorghum cultivated under *F. albida* is 1.7 times higher than those of plots without it [35].

3.3. Characteristics of Agricultural Production Systems in Fitri Watershed of Chad

In the Fitri watershed of Chad, different agricultural production systems are practised in Yao, Mochi and Zegue villages. The most promising are *Faidherbia albida* agroforestry parklands, mixed crops and Sorghum planted out (*Sorghum durra*) systems. The area allocated to each speculation varied from one village to another. Superficies dedicated to *Sorghum durra* are very low (3.70 ± 1.4 ha) (Figure 1).

The area of *F. albida* agroforestry parklands (7.32ha) and mixed crops systems (5.14ha) are large. The extension of surface areas of *F. albida* agroforestry parklands is due to the important role played: it increases the rate of water infiltration, improves soil fertility and conserves biodiversity by limiting the effects of climate changes. According to respondents, *Sorghum durra* planted out, on the contrary, degrades soil and produces poor yields. This situation results from the decrease in soil fertility and physical degradation of soils due to clearing practices and the destruction of plant cover. Thus, the presence of *F. albida* in agricultural production systems improves productivity (Figure 2).




Figure 1. Agricultural production systems in the watershed of Fitri.



Figure 2. *Ipomoea batatas* crop under *Faidherbia albida* trees.

F. albida is an important plant species used in soil restoration in the Sahelian zone [36]. It is a fixing nitrogen

species because of symbiotic association with *Rhizobium* spp. It uses this association to improve the soil fertility level in nitrogen in other to boost agricultural productivity. However, the dendrometric characteristics of trees found in these agricultural production systems of the watershed of Fitri reflect a certain degree of anthropization. Their density fluctuates between 34.42 in Mochi to 93.51 ± 2.7 stems /ha in Zegue (Table 3). The high density observed is registered in Zegue where woody species abounded and produced seeds which could permit their replenishment. This trend was reported in Burkina Faso [37] and Benin [38]. The diameter of trees varies from 0.32 ± 0.59 cm in Mochi to 0.72 ± 0.28 cm in Zegue. Despite the variability observed, there was no significant difference among the site ($0.7870 > 0.05$). These findings are similar to those reported in the Centre of Niger [39]. The presence of trees about 5 m in the height could be explained by the selective choice of the farmers.

Table 3. Dendrometric parameters of tree species.

Dendrometric parameters	Mochi	Yao	Zegue	Mean
Density (stems/ha)	34.42 ± 2.25	73.95 ± 3.48	93.51 ± 2.7	67.29 ± 2.81
Diameter (cm)	0.32 ± 0.59	0.47 ± 0.63	0.72 ± 0.28	0.50 ± 0.5
Height (m)	7.94 ± 2.9	10.16 ± 2.39	14.09 ± 5.1	10.73 ± 3.46

3.3.1. Ecological Importance of Species in Fitri Watershed in Chad

The ecological importance value index of the various species varies from 10.03 in *Acacia seyal* in the site of Zegue to 32.52 for *F. albida* in Moschi. Among the plant species recorded in Fitri watershed, four present index values equal to or more than 10% (Table 4). There is not a significant difference among the villages of Fitri watershed ($0.8714 > 0.05$). The importance value index of *F. albida* confirms what is above mentionned. It is an important ecological and socio-economic species: its leaves litter fertilize the soil, its tap roots increase the percentage of water infiltration and organic matter in the soil. Its fruits are excellent fodder for animals thus, *F. albida* plays an important role in enhancing resilience against climate changes in Fitri watershed of Chad.

Table 4. Percentage of importance value index of plant species.

Sites/Species	Mochi	Yao	Zegue	Mean
<i>Faidherbia albida</i>	32.52	23.25	15.62	23.80
<i>Balanites aegyptiaca</i>	28.84	20.31	17.16	22.10
<i>Prosopis juliflora</i>	16.07	21.58	13.16	16.94
<i>Mangifera indica</i>	12.63	11.19	10.36	11.39
<i>Tamarindus indica</i>	11.08	-	10.04	7.04
<i>Hyphaene thebaica</i>	12.56	-	11.14	7.90
<i>Borassus aethiopum</i>	-	13.11	11.06	8.06
<i>Acacia tortilis</i>	13.21	-	-	04.40
<i>Acacia senegal</i>	-	11.54	10.12	07.74
<i>Acacia seyal</i>	-	10.09	10.03	06.71
<i>Zizyphus mauritiana</i>	10.32	-	11.68	07.33

- = absent

Four of the species are present in all the villages of Fitri

watershed with an index value of importance $\geq 10.36\%$: *Faidherbia albida*, *Balanites aegyptiaca*, *Prosopis juliflora* and *Mangifera indica*. Among these tree species, *Mangifera indica* is an exotic one planted by farmers.

3.3.2. Ecological Importance of Families of the Species in the Watershed of Fitri

The analysis of the ecologically important families in Fitri watershed revealed that whatever the site, the main families

that exhibited Family Importance Index equal to or more than 15% are the following: Fabaceae, Balanitaceae, Mimosaceae and Arecaceae (Figure 3). The Fabaceae family is the top one with 32.04%. It regroups many species of great socio-economic and ecological importance. They are appreciated by farmers of the Fitri watershed. There is a significant difference among sites ($0.0134 < 0.05$).




Figure 3. Ecological families importance.

The steady management of families with high FIV needs to be considered in view to preserve the socio-economic and ecological services that they offer. In this way, the conservation of agroforestry species and promotion of agroforestry must be encouraged in Fitri watershed.

3.3.3. Ecological Service of Agricultural Production Systems

The carbon stock of Fitri watershed agricultural production

systems varies from 4.45 ± 0.9 in *Sorghum durra* fields established out to 39.41 ± 1.32 tC/ha in *F. albida* agroforestry parklands. The carbon sequestration potential of these systems oscillates between 16.1 ± 1.35 tCO₂eq/ha in *S. durra* established out and 144.64 ± 1.96 tCO₂eq/ha in *F. albida* agroforestry parklands. The total ecological service of the agricultural production systems of the watershed of Fitri of Chad is 201.41 t CO₂eq/ha (Table 5).

Table 5. Carbon stock accumulated by agricultural production systems.

Agricultural production systems	Biomass (t/ha)	Carbon (tC/ha)	Carbon (tCO ₂ eq/ha)
<i>F. albida</i> agroforestry parklands	82.13 ± 2.6	39.41 ± 1.32	144.64 ± 1.96
Mixed crops	23.1 ± 1.17	11.08 ± 1.9	40.67 ± 1.54
<i>Sorghum durra</i> planted out	9.28 ± 1.8	4.45 ± 0.9	16.1 ± 1.35
Total	123.51 ± 5.57	54.94 ± 4.12	201.41 ± 4.85

4. Discussion

4.1. Farmer Perceptions of Climate Changes

Climate changes are not perceived only by scientists, farmers are aware of their effects in the watershed of Fitri in Chad and have good pieces of knowledge on the variability in their villages through natural and physical factors such as yield decreasing, temperature growth, irregularity of precipitations, frequent floods, houses open to violent winds and dryness, etc. The high fluctuation of precipitations and temperatures registered in the Fitri watershed of Chad is

similar with the findings of many researchers who study the problematics of climate changes [40-43]. The rainy season gap has been put into evidence in subsaharian Africa by various authors [44, 45]. Concerning endogenous strategies developed by farmers, *Faidherbia albida* is a multipurpose tree species valued by farmers of the area (source of fodder for animals, restoration of soil fertility, medicines, fuelwoods and charcoal, etc.). A multifunctional tree species like this one is a «worker tree» which has many functions in the field [46]. In addition, diversification of crops was evoked by 2/3 of the respondents. Farmers who have at one disposal two kinds of cereal (*sorghum durra* and *Zea mays*) and one

leguminous crop (*Arachis hypogea* and/or *Vigna unguiculata*), are more resilient than others. Diversified crops can produce multiple yields in response to climate changes. The adoption of Sorghum established out has been cited by certain farmers. They reported that farmers who practice these crops of counter-season have a benefice of additional income which permits them to improve their living style.

4.2. Characteristics of Agricultural Production Systems in Fitri Watershed of Chad

Among the various agricultural systems, three linked to agroforestry systems present large areas (More than 3ha per Household). According to households, the ratio of crops associated with *F. albida* is bigger than that of the *S. durra* planted out in production systems. *F. albida* is a nitrogenous species that contributes to soil fertility. The species softens the light infiltration and its roots spread deep in the soil. These systems use litter and biomass easily decomposable permitting accommodation to the variability of climatic conditions. These findings are similar to those demonstrated in Kiki in Niger that fertility under *F. albida* is high. In the Sudano-sahelian zone of Cameroon, farmers claim that *Zea mays* growing under the species during the dry season withers latter than that which is far from it [46]. In addition, fruits of this species hold a high fodder value. They are particularly rich in the nitrogenous digestive matter [47]. Diversified Mixed crops reduce risks of crop yields and increase their ductibility [48].

For the flora in these systems, the following species present ecological importance: *Faidherbia albida* (23.80%), *Balanites aegyptiaca* (22.10%), *Prosopis africana* (16.94%) and *Mangifera indica* (11.39%). These plant species are important species of great socio-economic and ecological importance. They play an additional function of windbreak, by reducing the velocity of winds [49]. They said that they contribute to reinforcing resilience and mitigating climate changes. These findings are similar to those reported in Benin [50, 51] and Wofo (2008) in Cameroon [52]. Nevertheless, they are in another hand some findings which contrast with the previous in Northern Cameroon [53].

Concerning plant families, eight exhibited an important family index among which, the family of Fabaceae leads. Some studies in Africa reported the importance of this family in agroforestry systems [34, 54-56]. Meanwhile, the index of family importance is useful in the biodiversity conservation programm. Families with weak indexes are among the priorities for conservation [57] whereas those with high indexes to be well-managed and followed [58].

4.3. Ecological Service of Agricultural Production Systems in Fitri Watershed of Chad

The total ecological service of the agricultural production systems of Fitri watershed is 201.41t CO₂eq/ha. The two traditional agroforestry systems (*F. albida* agroforestry parklands, Mixed crops systems) store significant amounts of

carbon in their biomass. The ecological service of *F. albida* agroforestry parklands is 1.4 whereas that of *S. durra* planted out is 0.07 times compared to the total ecological service. The superiority of the *F. albida* agroforestry parklands in sequestering carbon in other systems was reported in West African Sahel [54]. These values concerning the carbon stock are higher than 24tC/ha reported in Parklands of Mali [59] and at the same time, fit in the interval (40-60 tC/ha) reported [60]. It is also higher than 5.046 tC/ha obtained in *Vitellaria paradoxa* agroforestry parklands in Cameroon [61]. Based on these findings, they can say that agroforestry systems are excellent carbon shafts.

5. Conclusion

Farmers have good knowledge of climate changes in their area. This study carried out in three villages of the watershed of Fitri permits to identify three agricultural production systems among which *F. albida* agroforestry parklands are the most important in increasing water infiltration, soil fertility and biodiversity conservation, thus limiting consequences of climate changes. Diversification of mixed crops reduces considerably risk of failure linked to climate changes. Ecological importance species are *Faidherbia albida*, *Balanites aegyptiaca*, *Prosopis juliflora* and *Mangifera indica*. Fabaceae, Balanitaceae, Mimosaceae, Arecaceae, Rhamnaceae, Capparaceae, Meliaceae and Asclepiadaceae are the main top families with an importance value index equal to or more than 10%. Plant diversity blooms better in Zegue than elsewhere in Fitri watershed. Agroforestry represents alternatives which can reverse actual trends of degraded natural ecosystems in the area. The high density of woody species shows that they abound in seeds which can permit their renewal. The development of an agroforestry system developed by farmers to enhance resilience against Climate changes could be encouraged in the broader context of sustainable development in the area of the watershed of Fitri of Chad. Agroforestry is an agro-ecological and climate-smart practice with huge potentials to reduce smallholder farmers' vulnerability and enhance their resilience faced with the predominantly negative effects of climate change. The next step will be focused on biodiversity conservation in the area.

Acknowledgements

This study received no funding. The authors thank the agriculturors of Fitri for their hard work and efficiency. They also greatly appreciate the valuable comments from the anonymous reviewers.

References

- [1] CILSS, 2016. Lutte contre les changements climatiques, les réalisations du CILSS, cobb. S (1982). Social support health through the life course, 42p.

- [2] Kiari Fougou H., 2014. Impacts des variations du niveau du lac Tchad sur les activités économiques des pêcheurs de la partie Nigérienne. Thèse de Doctorat de géographie, Université Abou-Moumoni, Niamey, Niger, 318p.
- [3] Moupeng B., 2006. Le lac Fitri: dynamique du système hydrologique quaternaire et actuel (Sahel tchadien). Thèse de Doctorat, Université de la Provence, Avignon, 171p.
- [4] Raimond C., Zakinet D., Mugele R., Kemsol N A., Mbagogo A., Yali-kunT., Brahim B. A., Madjigoto R., Schuster M., Sylvestre F., Deschamps P., 2017. Les nouveaux enjeux pour le lac Fitri, entre variabilité environnementale, croissance démographique et conflit d'usage. In: *livre des résumés étendus du colloque International Recherche croisée sur les écosystèmes lacustres tchadiens*, Ndjamen, Tchad, 25-27 avril 2017, 601p.
- [5] Scheffers A. M., Scheffers S. R., Kelletat D. H., 2012. The coastline of the world with google Earth: undertaking our environment Coastal Research Library 2, springer, Dordrecht, 293p.
- [6] Schuster M., 2002. Sédimentologie et paléontologie des séries à vertébrés du paléo lac Tchad depuis le Miocène supérieur. Thèse de Doctorat, Université Louis Pasteur de Strasbourg, 152p.
- [7] Sandjong Sani R. C., Ntoupka M., Adamou I et Vroumsia T., 2013. Etude écologique du Parc National de Mozogo-Gokoro (Cameroun): prospection préliminaire de la flore ligneuse et du sol pour la conservation et son aménagement, In. *I. Biol. Chem. Sc.* 7 (6): 2434-2449.
- [8] Marty., Zakinet D., Khamis D., Bernard C., 2012. Analyse de l'évolution du Fitri. *Document principal*. République du Tchad, programme d'hydraulique pastoral au Tchad central phase 2, Antea-Iram, 128p.
- [9] Young A., 1989. Agroforestry for soil conservation. Wallingford, oxon, u. k: CAB. International., *International Concl for Research in Agroforestry*, 23-52.
- [10] Mapongmetsem P. M., Nduryang J. B. N., Fawa G., Dona A., 2015. Contribution à la connaissance des produits forestiers non ligneux de la zone soudano-sahélienne du Cameroun. Kapseu C., Nzié W., Nso E., Silechi J., Ngomo H., (Eds). Biodiversité et changements globaux: valorisation des effluents des industries, des résidus agro-pastoraux et forestiers. *Actes du Colloque International* du 21 au 23 juillet 2015 à Ngaoundéré, 139-147.
- [11] Nair P. K. R. & Garrity D., 2012. Agroforestry; the future of global land use, advance in agroforestry *springer*. Dordrecht, 541p.
- [12] Mapongmetsem P. M., Dona A., 2010. Land use systems and biodiversity conservation east Tandjile, Chad. *Scripta Bot. Belg*, 46: 285.
- [13] Dona A., Mapongmetsem P. M., Dongock N. D., Pamboudem N. A., Fawa G. and Aoudou D. S., 2016. Phytodiversity and carbon stock in Sudanian savannahs zone of Tandjile-East of Chad, *International Journal of Applied Research*, 2 (9): 455-460.
- [14] Torquebiau E., Mary F & Sibelet N., 2002. Les associations agroforestières et leurs multiples enjeux. *Bois et forêts des Tropiques*, 271: 23-34.
- [15] Garba I., Touré I., Ickowicz A., 2012. « Evolution historique de la pluviosité ». In système d'information sur le pastoralisme au Sahel. Atlas des évolutions des systèmes pastoraux au Sahel 1970-2012. *Fao et Cirad*, ed: 8-11.
- [16] Woodfine A., 2008. L'adaptation au changement climatique et l'atténuation de ses effets en Afrique Subsaharienne au moyen des pratiques de gestion durable des terres, *Terrafrica*, 90p.
- [17] Kemsol Nagorngar A., 2018. Dynamique des cultures de décrue dans les zones lacustres Soudano-Sahéliennes de 1985 à 2015: cas des lacs Fitri et Iro au Tchad. Thèse de doctorat, Université Felix-Houphouët-Boigny d'Abidjan-Cocody, 178p.
- [18] Higgings I. S., Shackleton M. C., Robinson R. E., 1999. Changes in woody community structure and composition under contrasting lean use systems in semi-arid Savanna. South Afrique, *Journal of biogeography*, 26: 2019 – 629.
- [19] RGPH2, 2009. Deuxième recensement général de la population et de l'habitat. Ministère du plan, Tchad, 163p.
- [20] P-SIDRAT, 2013. Schéma régional d'aménagement du territoire (SRAT) du Batha. Rapport final, consortium Agrer-Image, Matuh, 49p.
- [21] Ngom D., 2013. Diversité végétale et quantification des services écosystémiques de biosphère du Ferlo (Nord-Sénégal). Thèse, ED-SEV/UCAD, Dakar, 167p.
- [22] Yédomonhan H., Hounadagba C. J., Akoéginou A., Vander Maesen L. J. G., 2008. Structure et diversité floristique de la végétation des inselbergs du secteur méridional du centre-Bénin. *Syst. Geogr. Pl*, 78: 111- 125.
- [23] Felfili J. M., Silva Junior M. C., Sevilha A. C., Fagg C. W., Walter B. M. T., Nogueira P. E & Rezende A. G., 2004. Diversity floristics and structural patterns of cerrado vegetation in Central Brazil, *Plant Ecology*, 175: 37 – 48.
- [24] Neelo J., Teketay D., Kashe K., Masamba W., 2015. Stand structure, diversity and regeneration statut of woody species in open and exclosed. Dry woodland sites around Malapo farming areas of the Okavango Delta, Northeastern Bostwana. *Open Journal of Forestry*, 5: 313 – 328.
- [25] Nusbaumer L., Gautier L., Chatelain C&Spichiger R., 2005. Structure et composition floristique de la forêt classée du Scio (Côte d'Ivoire). Etude descriptive et comparative. *Candollea*, 60: 394-443.
- [26] Gonmadje, C. F., Doumenge, C., McKey, D., Tchouto, M. G. P., Sunderland, T. C. H., Balinga, M. P. B., Sonké, B., 2011. Tree diversity and conservation value of the Ngovayang massif, Cameroon. *Biodiversity and Conservation*, 20: 2627-2648.
- [27] Gonmadje, C. F., Doumenge, C., Sunderland, T. C. H., Balinga, M. P. B., Sonké, B., 2012. Analyse phytogéographique des forêts d'Afrique Centrale: le cas du massif de Ngovayang (Cameroun). *Plant Ecology and Evolution*, 145 (2): 152-164.
- [28] Kacholi D. S., 2014. Analysis of structure and Diversity of the Kilengwe Forest in the Morogoro Region, *Tanzania International Journal of Biodiversity*. Volume 2014, Article ID 516840, 8p.
- [29] MBow C., Verstraete M and Neufeldth, 2014. Allometric models for aboveground biomass in dry savanna trees of the Sudan and Sudan-Guinean ecosystems of southern Senegal, pp. 340-347.

- [30] GIEC, 2007. Résumé à l'intention des décideurs. Changements climatiques 2007: Les éléments scientifiques. Contribution du Groupe de travail I au quatrième rapport d'évaluation du Groupe d'experts intergouvernemental sur l'évolution du climat, Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K. B. Averyt, M. Tignor et H. L. Miller (eds.). Cambridge University Press, Cambridge, UK et New York, NY, USA. 2007.
- [31] GIEC, 2006. Guide pour l'inventaire national des gaz à effet de serre agriculture, foresterie et autre usage des terres. Institute for Global Environmental Strategies, Japon, 4: 46-52.
- [32] Valentini G., 2007. Évaluation de la séquestration du carbone dans les plantations agroforestières et des jachères issues d'une agriculture migratoire dans les territoires autochtones de Talamanca au Costa Rica. Mémoire de MSC en Biologie végétale, Université de Laval Québec, Canada, 128p.
- [33] Roupsard O., Ferhi A and Granier A., 2002. Reverse phenology and dry-season water uptake by *Faidherbia albida* (Del.) A. Chev. in an agroforestry parkland of Sudanese West Africa, *Functional Ecology*, 13 (4): 460 – 472.
- [34] Gassama –Dia Y. K., San'Ed and N'Doye M., 2003. Reproductive biology of *Faidherbia albida* (Del.) A. Chev. *Silva Fennica*, 37: 429-436.
- [35] Gonzalez P., Tucker C. J., 2012. Tree density and species decline in the African Sahel attributable to climate, *Journal of Arid Environment*, 78: 55-64.
- [36] Cesse M., Bationo B. A., Traore S et Bourssim I. J., 2018. Perception d'espèces agroforestières et de leurs services écosystémiques par trois (3) groupes ethniques du bassin versant de BOURA, zone Soudanienne du Burkina-Faso. *Bois et Forêts de Tropiques*, 338: 29-42.
- [37] Yameogo G., Yelemou B., Boussim I. J., Traoré D., 2013. Gestion du parc agroforestier du terroir de VIPALOGO (Burkina-Faso): contribution des ligneux à la satisfaction des besoins des populations. *Int. J. Bio. Chem. Sci.*, 7 (3): 1087-1105.
- [38] Biao S. S. H., Natta A. K., Dicko A et Kouagou M'M., 2016. Typologies des systems agroforestiers et leurs impacts sur la satisfaction des besoins des populations rurales au Benin, 14p, web <http://www.slire.net> & <http://www.inrab.org>
- [39] Abourhamane H., Morou B., Rabiou H et Mahamane A., 2013. Caractéristiques floristiques, diversité et structure de la végétation ligneuse dans le Centre-Sud du Niger: cas du complexe des forêts de Dan Kada Dodo-Dan Gado. *Int. J. Biol. Chem. Sci.*, 7 (3): 1048-1068.
- [40] Kanohin F., Saley M. B & Savane L., 2009. *European Journal of Scientific Research*, 26: 209-222.
- [41] Bambara D., Bilgo A., Lompo F., Hien V., 2011. *Int. J. Biol. Chem. Sci.*, 5 (6): 2415 – 2433.
- [42] Mamah M., Baye-Niwah C., Hamawa Y., Dangai Y., Oumarou Z., Abdoulaye H., Mal Ali, Fawa G. & Mapongmetsem P. M., 2019. *Borassus aethiopum* Mart. agroforestry parklands and climate change mitigation in Cameroon. *International Journal of Agriculture and Environmental Research* 5 (4): 436-455. www.ijaer.in
- [43] Dangai Y., Hamawa Y., Tsobou R., H. Z. Oumarou H. Z., Mapongmetsem P. M., 2021. Carbon stocks in *Daniellia oliveri* agroforestry parklands in the Sudano - sahelian zone of Cameroon. *Environmental Challenges*, 5: 1-8. Journal homepage: www.elsevier.com/locate/envc
- [44] Marema T & Growley J., 2004. Impact des changements environnementaux sur la migration humaine, études de cas: Sénégal et Côte d'Ivoire, juin, UNESCO, 29 p.
- [45] Salack S., Traoré B & Sarr B., 2006. Synthèse sur les collecte, la mise en forme et le stockage des données climatologiques des pays du CLSS et étude d'impacts des changements climatiques sur la production agricole au Sahel, Rapport de stage, Centre AGRHYMET, 95p.
- [46] Smith Dumont E, Bonhomme S., Fergus S., 2015. Guide technique d'agroforesterie pour la selection et la gestion des arbres au NORD-KIVU. RDC, WAC. 130p.
- [47] Bationo B. A., Kalinganiré A. et Bayala J., 2012. Potentialités des ligneux dans la pratique de l'agriculture de conservation dans les zones arides et semi-arides de l'Afrique de l'Ouest: Aperçu de quelques systèmes candidats. ICRAF Technical Manual no 17 Nairobi: *World Agroforestry Centre*, 12: 87-94.
- [48] Alcade C. S., Achigan-Dako E. G., Orou G. G and Ahanchédé A., 2015. Farmer's knowledge and perception of diversified farming systems in Sub-Humid and Semi-Arid Areas in Benin, *Sustainability*, 7, 6573-6592.
- [49] Sobola I., Aubricot O., Puschiasis O., Duplan T., Grimald J., Hugonnet M et Buchheit P., 2015b. Changement climatique et ressource en eau en Himalaya: enquête auprès des villageois dans quatre unités géographiques du bassin de la Koshi, Nepal. *Revue de géographie Alpine*, 103 (2), 1-25.
- [50] Boureima M., Sissoko K., Zougmore R., Dieye K., Amadou M., Moussa A. S., Foerch W., Garlick C., Ochieng S., Kristjanson P., Thornton P. K., 2011. Résultats des enquêtes de base des ménages du site de Kolle, Niger, 17p.
- [51] Adamou C. A., Mana A., Missikpode R., Sinsin R., 2009. Cartographie et caractérisation floristique de la forêt marécageuse de LOKOLI (Benin), 14p.
- [52] Wofu G., 2008. Les aires protégées de l'Extrême Nord du Cameroun entre politique et conservation et pratiques locales. Thèse de Doctorat en géographie Aménagement-Environnement, Université d'Orléans, 325p.
- [53] Awono Mvondo J. P., Boukong A., Mainam F., Yombo G., Njokou Tchoutang G., Beyegue-Djonko H., Fertilisation des sols dans les monts Mandara à l'Extrême-Nord du Cameroun: du diagnostic aux recommandations. 8 p. fihal-00142746f
- [54] Asako Takimoto, Ramachandran Nair P. K., Vimala D. Nair, 2008. Carbon stock and sequestration potential of traditional and improved agroforestry systems in the West African Sahel. *Agriculture, Ecosystems and Environment*, 125: 159-166.
- [55] Mama A., Sinsin B., Cannière C & Bogaert J., 2013. Anthropisation et dynamisme des paysages en zone Soudanienne au Nord du Bénin, in *Tropicultura*, 3: 78-88.
- [56] Barmos S., Amani A., Soumana I., Ichaou A., Karim S & Mahamane A., 2019. Structure et diversité des parcs agroforestiers adjacents à la forêt protégée de Bahan Rafi, Niger-Afrique de l'Ouest. *Afrique Science*, 15 (2): 166-185, <http://www.afriquescience.net>
- [57] Konan D., Bakayoko A., Trabi F. H., Bitignon B. G. A & Piba S. C., 2015. Dynamisme de la structure diamétrique du peuplement ligneux des différents biotopes de la forêt classée de Yapo-Abbé, Sud de la Côte d'Ivoire. *Journal of Applied Biosciences*, 94: 8869 – 8879.

- [58] Gueulou N., Ouattara N. D., Konan D., Gnahore E., Missa K & Bakayako A., 2018. Diversité floristique et structurale de la forêt galerie du Bandama dans la Réserve Scientifique de Lamto en Côte d'Ivoire. *Afrique SCIENCE*, 14 (4): 439 -452.
- [59] Traoré K., Ganry F., Olivier R., Gigou R., 2004. Litter production and soil fertility in a *Vitellaria paradoxa* parkland in a Catena in Southern Mali. *Arid land Res. Manag*, 18 (4): 359- 368.
- [60] Palm C. A., Woomer P. L., Alegre J., Arevalo L., Castella C., Cordeiro D. G., Feigl B., Hairiah K., Kotto-Same J., Mendes A., Moukam A., Murdiyarso D., Njomgang R., Parton W. J., Ricse A., Rodrigue V., Sitompul S. M., Van Noordwij K. M., 2000. Carbon sequestration and trace gaz emissions in slash-and-burn and alternative land uses in the humid tropics. Final Report, Alternative to slash and Burn (ABS). Climate change working Group, *phase III ICRAF*, Nairobi, Kenya, 29 p.
- [61] Peltier R., Njiti F. C., Ntoupka M., Manlay R., Henry M., Morillon V., 2007. Evaluation du stock de carbone et productivité en bois d'un parc à karité du Nord-Cameroun. *Revue Bois des Tropiques*, 294 (4): 39 – 50.